

---

## LMBP BACTERIAL HOST STRAIN

## MG1655 seqA-eYFP

These validated data are a snapshot at a given moment; further updates are always possible.

---

<u>Species:</u>	<i>Escherichia coli</i>
<u>Group:</u>	K12
<u>Accession number:</u>	<b>LMBP 9776</b>
<u>Deposit date:</u>	11/01/2016
<u>Depositor:</u>	Prof. Dr J. Michiels <sup>1</sup> (constructed by T. Swings <sup>1</sup> ) <sup>1</sup> Centre of Microbial and Plant Genetics, Department of Microbial and Molecular Systems, KU Leuven, Leuven, Belgium
<u>Other culture collection numbers:</u>	/
<u>Containment level:</u>	This strain has been assigned the containment level 'Class 1' following the European Directive 2009/41/EC on the contained use of genetically modified organisms, and its updates (see also the <a href="#">Belgian risk group classification</a> ).
<u>Medium:</u>	LB-Miller
<u>Selection marker:</u>	Kanamycin (50 µg/ml)
<u>Cultivation temperature:</u>	37°C
<u>Original reference:</u>	Mika et al., Faraday Discuss. 184 (2015), 425-450 [PMID: <a href="#">26449690</a> ]
<u>Related reference:</u>	/
<u>Genotype*:</u>	<i>F</i> λ <sup>-</sup> <i>ilvG</i> <sup>-</sup> <i>rfb-50 rph-1 seqA-eYFP:km</i> <sup>R</sup>
<u>Phenotype:</u>	Km <sup>R</sup>
<u>Properties:</u>	The native <i>seqA</i> gene was fused at its 3' end to the fluorescent protein gene <i>eYFP</i> , separated with a linker (ASPPPGRSR) and followed by a kanamycin resistance marker. The entire cassette was integrated in the chromosome. This strain can be used to study the subcellular localization of the <i>E. coli</i> DNA binding protein SeqA.
<u>Restricted use:</u>	- BCCM MTA - The depositor will be informed of the customer's identity upon release of a sample outside the depositor's department or outside the departments in which BCCM/LMBP is embedded, namely UGent-DBMB and VIB-IRC.

\* Source: [http://openwetware.org/wiki/E.\\_coli\\_genotypes#MG1655](http://openwetware.org/wiki/E._coli_genotypes#MG1655) + Mika et al. (2015) [PMID: [26449690](#)]

### **Culture recovery and preservation instructions**

The enclosed culture has been grown overnight to saturation, confirming its viability. BCCM/LMBP advises to recover it immediately on receipt before use or storage.

Recovery: subculturing into liquid or solid medium according to the cultivation conditions described above.

Long-term preservation: lyophilisation of the subculture  
cryopreservation (at -80 °C at the least)