
LMBP BACTERIAL HOST STRAIN

K12 Δ H1 Δ trp

<u>Species:</u>	<i>Escherichia coli</i>
<u>Group:</u>	K12
<u>Accession number:</u>	LMBP 69
<u>Deposit date:</u>	01/01/1998
<u>Depositor:</u>	Prof. E. Remaut ^{1 2} ¹ Department for Molecular Biomedical Research, VIB, Ghent, Belgium ² Department of Molecular Biology, Ghent University, Ghent, Belgium
<u>Medium:</u>	LB
<u>Selection marker:</u>	/
<u>Cultivation temperature:</u>	28°C
<u>Original reference:</u>	Bernard et al., Gene 5 (1979), 59-76 [PMID: 372049]
<u>Related reference:</u>	Remaut et al., Gene 15 (1981), 81-93 [PMID: 6271633]
<u>Genotype:</u>	<i>lacZam</i> Δ (<i>bio-uvrB</i>) Δ (<i>trpE-A</i>) ² <i>rpsL</i> λ (<i>Nam7 Nam53 cI857</i> Δ H1)
<u>Phenotype:</u>	Sm ^R
<u>Properties:</u>	Expression host for plasmids containing phage λ 's PL or PR promoter. Induction is obtained by shifting the culture from 28°C to 42°C. The resident λ is a defective lysogen. Δ H1 removes part of <i>cro</i> and all genes to the right of it. <i>cI857</i> is ind ⁻ .
<u>Restricted use:</u>	BCCM MTA

Culture recovery and preservation instructions

The enclosed culture has been grown overnight to saturation, confirming its viability. BCCM/LMBP advises to recover it immediately on receipt before use or storage.

Recovery: subculturing into liquid or solid medium according to the cultivation conditions described above.

Long-term preservation: lyophilisation of the subculture
cryopreservation (at -80 °C at the least)